

Conscientious Objection in Medicine: Psychological consequences in women who face Legal Termination of Pregnancy in the Region of Murcia

BACKGROUND

- Perinatal loss is a stressful situation in women's life (and their families) (Daugirdaité, van den Akker & Purewal, 2015)¹
- Losses due to legal termination by medical criteria carry several traumatic moments

Aim

To describe women experience and to explore symptoms of Posttraumatic Stress Disorder (PTSD) in relation to the medical care received during the pregnancy termination

METHOD

Instrument: On-line ad hoc questionnaire + PCL-6 (symptoms of PTSD) (Milles, Marchall & Schell, 2008)²

Data collection: Between February and August 2019

Participants: 27 women. Age average = 33,44 years-old (s. d. = 3,68; range = 28-43). 22,22% Region of Murcia

RESULTS

• Weeks pregnancy average

17,33 (4,83) weeks + 3,45 days; range= 8-32

• Decision making date

18,41(4,63)weeks+4,63 days; range=13-32

• Interruption date

19,19(4,84) weeks +4,06 days

• Interruption in Reference Center


18,51%. (Murcia=0%)

• Prejudice by medical staff

14,87% (Murcia=33,33%)

	Support	Reference Center	Private Clinic
Economical	18,51% Satisfaction=1,59/5 (Murcia = 33,33%;1/5)	0%	
Bureaucratic	33,33% Satisfaction = 3,5/5 (Murcia = 0%).	13,79% Satisfaction = 4,2/5	
Psychological	18,51% (1 accepted) Satisfaction = 5/5 (Murcia = 0%)	3,44 % Satisfaction = 1/5 (Murcia = 0%).	

(Satisfaction: 1=very poor - 5= excellent)



PCL-6 → 75,86% ≥14 positive symptoms

Average: ■ 22,89 □ 22,42 ■ 23,89

MCsatisfaction * PCL-6 correlation:

r = -0,63;p<0,001

CONCLUSIONS

- Perinatal loss is a bad experience in addition due to the medical care. PTSD symptoms are present and correlates with low Medical Care Satisfaction
- Women from Region of Murcia showed worst Medical Care Satisfaction.
- Psychological professional support is required.
- Medical standards need revision and further research must be conducted.